

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Belimumab (Benlysta®)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:/ Referral Status: New Referral Dose or Frequency Change Renewal				
Patient Name: Patient Address: Patient Phone Number: Date of Birth:/ Weight:kg	Primary Insurance: Member ID: Secondary Insurance: Member ID:			
Diagnosis ☐ Drug-induced systemic lupus erythematosus (M32.0) ☐ Systemic lupus erythematosus organ or system involvement unspecified (M32.10) ☐ Endocarditis in systemic lupus erythematosus (M32.11) ☐ Pericarditis in systemic lupus erythematosus (M32.12) ☐ Lung involvement in systemic lupus erythematosus (M32.13) ☐ Glomerular disease in systemic lupus erythematosus (M32.14) ☐ Tubulo-interstitial nephropathy in systemic lupus erythematosus (M32.15) ☐ Other organ or system involvement in systemic lupus erythematosus (M32.19) ☐ Other forms of systemic lupus erythematosus (M32.8) ☐ Systemic lupus erythematosus, unspecified (M32.9) ☐ Other Diagnosis Code (ICD-10): ☐ Other Indication: ☐ Target start date: ☐ Target start date:	Labs Frequency CMP CBC w/diff Every infusion CBC w/o diff Other:			
Previously tried and failed therapies (include dates):				
Belimumab (Benlysta) Dose 10 mg/kg mg/kg mg/kg mg	Frequency Induction Maintenance ☐ On weeks 0, 2, 4 ☐ Every 4 weeks ☐ Every weeks Date of last infusion://			



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Pre-Medications	☐ Acetaminophen 650 mg PO once		Hydration	
and Pre-Protocol	☐ Loratadine 10 mg PO once		□ LR	
(ordered at	☐ Diphenhydramine once		☐ Sodium Chloride 0.9%	
physician discretion)	□25 mg □50 mg		☐ Other	
	□ PO □IV			
	☐ Famotidine 20 mg IV once		mL atmL/hr	
	☐ Hydrocortisone 100 mg IV once		☐ Before ☐ During ☐ Following	
	☐ Methylprednisolone 125 mg IVP once			
	☐ Other:			
	a other.			
Flushing Protocol	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL	□ Не	parinUnits/mLmL	
(pre and post	as needed for line care	as nee	eded for line care	
medication)				
Hypersensitivity	Sodium chloride 0.9% bolus 500 mL onc	e as nee	ded for hypotensive management (SBP <90mmHg)	
Panel • Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches,				
Together Care	ogether Care generalized pain, back pain, abdominal cramping			
Hypersensitivity	• Albuterol 2.5 mg/3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia,			
Panel will be	bronchospasm, wheezing, dyspnea, for 2 doses			
ordered to provide	Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea			
emergency	• Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to			
supportive care	moderate anaphylaxis for 3 doses			
medication therapy	• Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion			
as necessary.	reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea,			
	cognitive changes, generalized rash, chest pain/pressure			
	Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate			
	hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors,			
	localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping,			
	uneasiness, agitation, feeling of impending doom			
	Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe When the second severe as the second second severe as the second			
	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or			
tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure				
Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hydrocortisone reaction (grade 3) including systelia RD 90 00 mg IVs bradycardia or				
hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure				
	tacifycardia, ffypoxerilla, dyspirea, cogilii	tive citai	iges, generalized rash, chest pani, pressure	
Provider Name:		Provid	ler Signature:	
Attanding Physician N	lama	Drovio	lor NDI	
Acceliaing Physiciali N	lame:	FIUVIC	ler NPI:	
Office Phone Number	:	Office	Fax Number:	
(If ordering provider is an advanced practice practitioner)				
		uro		
Note: This order is valid for 12 months from date of physician signature.				