

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Efgartigimod alpha-fcab (Vyvgart®)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:// Referral Status: □ New Referral □ Dose or Frequency Change □ Renewal					
Patient Name: Patient Address: Patient Phone Number: Date of Birth: Weight: kg Height: Allergies:		Primary Insurance: Member ID: Secondary Insurance: Member ID:			
Diagnosis ☐ Myasthenia gravis (G70.0) ☐ Myasthenia gravis without (acute) exacerbation (G70.00) ☐ Myasthenia gravis with (acute) exacerbation (G70.01) ☐ Other Diagnosis Code (ICD-10):		□ СВС	C w/diff C w/o diff er:	Frequency □ Every infusion □ Other:	
Efgartigimod alpha-fcab (Vyvgart®) 10 mg/kg in 0.9% sodium chloride (total volume 125 mL) infused over 1 hour. Nursing note: Administer with 0.2 micron in-line filter. Monitor patient for 1 hour post-administration for any signs and symptoms of hypersensitivity reactions.					
Dose ☐ 10 mg/kg (maximum dose: 1,200 mg) ☐ mg/kg ☐ mg		Frequency □ Every 1 week □ Every weeks Date of last infusion://			
Pre-Medications and Pre-Protocol (ordered at physician discretion)	□ Acetaminophen 650 mg PO once □ Loratadine 10 mg PO once □ Diphenhydramine once □ 25 mg □ 50 mg □ PO □ IV □ Famotidine 20 mg IV once □ Hydrocortisone 100 mg IV once □ Methylprednisolone 125 mg IVP once □ Other:		☐ LR ☐ Sodium Chl ☐ Other	Hydration loride 0.9% mL at mL/hr During	



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Flushing Protocol (pre and post	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL as needed for line care	☐ HeparinUnits/mLmL as needed for line care		
medication) Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy as necessary	 Acetaminophen 650 mg once as needed generalized pain, back pain, abdominal of Albuterol 2.5 mg /3 mL (0.083%) nebuliz bronchospasm, wheezing, dyspnea, for 2 Albuterol HFA inhaler 2 puffs q4hs as needed to Epinephrine injection 0.3 mg IM every 1 moderate anaphylaxis for 3 doses Famotidine injection 20 mg IV over 2 min reaction including systolic BP 80-90 mm and cognitive changes, generalized rash, cheeded to Diphenhydramine injection 25 mg IV over hypersensitivity/infusion reaction (grade localized rash/hives, vomiting, nausea, puneasiness, agitation, feeling of impendiction Diphenhydramine injection 50 mg IV over hypersensitivity/infusion reaction (grade tachycardia, hypoxemia, dyspnea, cognite hypersensitivity/infusion reaction (grade tachycardia) 	ter solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, 2 doses seded for hypoxemia, bronchospasm, wheezing, dyspnea 5 minutes as needed for SBP less than 90mmHg, mild to nutes once as needed for severe hypersensitivity/infusion Hg, bradycardia or tachycardia, hypoxemia, dyspnea, st pain/pressure er 1 minute once as needed, moderate 2) including temperature greater than 100.5, rigors, bruritis, flushing, dizziness, back pain, abdominal cramping, and doom er 1 minute once as needed for severe 2 3), including systolic BP 80-90 mmHg, bradycardia or tive changes, generalized rash, chest pain/pressure		
Provider Name:		Provider Signature:		
Attending Physician Name:		Provider NPI:		
Office Phone Number:		Office Fax Number:		
(If ordering provider is an advanced practice practitioner) Note: This order is valid for 12 months from date of physician signature.				