

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Infliximab (Remicaid ®) or Biosimilar

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: /						
Referral Status: ☐ New Referral ☐ Dose or Frequency Change ☐ Renewal						
Patient Name: Patient Address: Patient Phone Number: Date of Birth:/ Weight:kg	Primary Insurance: Member ID: Secondary Insurance: Member ID:					
☐ Rheumatoid Arthritis (M06) specified joint and	Labs □ BMP	Frequency				
laterality ICD 10:		☐ Every infusion				
□ Ankylosing Spondylitis (M45)	☐ Hepatic Panel	Other:				
☐ Psoriatic Arthropathy (L40.59)	☐ CBC w/diff					
☐ Regional Enteritis Unspecified (K50.90)	☐ CBC w/o diff					
☐ Ulcerative Enterocolitis (K51.00)	☐ CRP					
☐ Ulcerative Colitis Unspecified (K51.90)	☐ ESR					
☐ Other Diagnosis Code (ICD-10):	☐ Other:					
☐ Other Indication:						
Target start date:						
Previously tried and failed therapies (include dates):	I					
Infliximab or Biosimilar □ Pharmacy to Select* □ Renflexis® (infliximab-abda) - preferred □ Inflectra® (infliximab-dyyb) □ Remicaid® (infliximab) *Pharmacist will work with financial coordinator to select product based on patients and patients are producted based on patients and product based on patients and patients are producted based on patients and patients are producted based on patients and patients are producted based on patients are producted by produc						
Dose	Frequency					
☐ 3 mg/kg	Induction	Maintenance				
□ 5 mg/kg	\square weeks 0, 2 and 6	☐ Every 6 weeks				
□ 7.5 mg/kg □ 10 mg/kg		☐ Every 8 weeks				
□ 10 mg/kg □ mg/kg		☐ Every weeks				
mg	Date of last infusion:/					



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Pre-Medications	☐ Acetaminophen 650 mg PO once		Hydration			
and Pre-Protocol	☐ Loratadine 10 mg PO once		□ LR			
(ordered at	☐ Diphenhydramine once		☐ Sodium Chloride 0.9%			
physician discretion)	□25 mg □50 mg		☐ Other			
	□ PO □IV					
	☐ Famotidine 20 mg IV once			mL at	mL/hr	
	☐ Hydrocortisone 100 mg IV once		☐ Before	\square During	☐ Following	
	☐ Methylprednisolone 125 mg IVP once					
	☐ Other:					
Flushing Protocol	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL	□ Не	parin	Units/mL	mL	
(pre and post	as needed for line care	as nee	eded for line car	re		
medication)						
Hypersensitivity	• Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg)					
• Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches,						
Together Care	generalized pain, back pain, abdominal c		_			
Hypersensitivity	• Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia,					
Panel will be	bronchospasm, wheezing, dyspnea, for 2 doses					
ordered to provide	Albuterol HFA inhaler 2 puffs q4hs as ne			-		
emergency	• Epinephrine injection 0.3 mg IM every 1.	5 minut	es as needed fo	or SBP less than 9	OmmHg, mild to	
supportive care	moderate anaphylaxis for 3 doses			. ,		
medication therapy	Famotidine injection 20 mg IV over 2 min					
if necessary	reaction including systolic BP 80-90 mml	_	-	ycardia, nypoxer	nia, dyspnea,	
	 cognitive changes, generalized rash, chest pain/pressure Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate 					
	hypersensitivity/infusion reaction (grade				100 E rigors	
	localized rash/hives, vomiting, nausea, p	-		_	_	
				ess, back pairi, ab	dominal cramping,	
	 uneasiness, agitation, feeling of impending doom Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe 					
	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or					
	tachycardia, hypoxemia, dyspnea, cognit	-		_	-	
	Hydrocortisone sodium succinate injecti					
	hypersensitivity/infusion reaction (grade					
tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure						
Provider Name:		Provid	ler Signature: _			
Attending Physician Name: Provide		ler NPI:				
Office Phone Number: Office		Fax Number:				
(If ordering provider is an advanced practice practitioner)						
Note: This order is valid for 12 months from date of physician signature.						
Note: This order is valid for 12 months from date of physician signature.						