

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Iron Infusion

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:// Referral Status: □ New Referral □ Dose or Frequency Change □ Renewal				
Referral Status. In New Referral Dose of Frequency Change In Reflewar				
Patient Name:	Primary Insurance:			
Patient Address:	Member ID:			
Patient Phone Number:	Secondary Insurance:			
Date of Birth:/	Member ID:			
Weight:kg Height:cm				
Allergies:				
Diagnosis	Labs			
☐ Iron Deficiency Anemia (D50.9)	□ CBC			
☐ Other Diagnosis Code (ICD-10):	☐ Iron Studies (Iron, T-sat, TIBC, Ferritin)			
Is the patient on hemodialysis? Yes No	□ Phosphorus			
Did the patient have an inadequate response to oral iron	☐ Other:			
supplements? \square Yes \square No				
Target start date:	Hemoglobin: Date:			
Talget start date:	Ferritin: Date:			
Iron Produc	ct Selection			
☐ Pharmacist to	select and dose			
Ferumoxytol (Feraheme) – TH Tier 1 Preferred Therapy	Iron Sucrose (Venofer) – TH Tier 1			
\square 1020 mg IV over 60 minutes x once	☐ 100 mg IV push every 4 weeks			
☐ 510 mg IV over 30 minutes weekly x 2 doses	☐ 200 mg IV push 3 times weekly x 5 doses			
☐ Pharmacist to dose	☐ 200 mg IV push weekly x 5 doses			
☐ Other:	☐ 300 mg IV infusion every 2 weeks x 2 doses			
	-followed by-			
Sodium Ferric Gluconate (Ferrlecit) – TH Tier 1	400 mg IV infusion x 1 dose			
125 mg IV infusion 3 times weekly x 8 doses	☐ Pharmacist to dose			
Pharmacist to dose	☐ Other:			
☐ Other:	Dose mg Sig Total # of Doses			
Iron Dextran (Infed) – TH Tier 1	Ferric Carboxymaltose (Injectafer) – TH Tier 2			
☐ 25 mg IV infusion test dose	Non-preferred − must answer one of the first two boxes in addition to selecting dose ☐ Intolerance to other IV iron products			
-followed by-	-OR-			
975 mg IV infusion x once	☐ Insurance authorization requires use for treatment			
☐ 1000 mg IV infusion once (ONLY if tolerated previously)	-AND-			
☐ Pharmacist to dose	☐ 750 mg IV push weekly x 2 doses			
Other:	\Box 15 mg/kg IV push weekly x 2 doses (if < 50 kg)			
Administration time: \Box 1-hour infusion \Box 4-hour infusion	☐ Pharmacist to dose			



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Pre-Medications	lications		Hydration		
and Pre-Protocol	☐ Loratadine 10 mg PO once		□ LR		
(ordered at	☐ Diphenhydramine once		☐ Sodium Chloride 0.9%		
physician discretion)	□ 25 mg □ 50 mg		☐ Other		
	□ PO □IV				
	☐ Famotidine 20 mg IV once		mL atmL/hr		
	☐ Hydrocortisone 100 mg IV once		☐ Before ☐ During ☐ Following		
	☐ Methylprednisolone 125 mg IVP once				
	☐ Other:				
	Dottier.				
Flushing Protocol	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL	☐ He	parinUnits/mLmL		
(pre and post	as needed for line care	as nee	eded for line care		
medication)					
Hypersensitivity	Sodium chloride 0.9% bolus 500 mL once	e as nee	ded for hypotensive management (SBP <90mmHg)		
Panel • Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches,					
Together Care					
Hypersensitivity	• Albuterol 2.5 mg/3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia,				
Panel will be	bronchospasm, wheezing, dyspnea, for 2 doses				
ordered to provide	• • • • • • • • • • • • • • • • • • • •				
emergency	• Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to				
supportive care	moderate anaphylaxis for 3 doses				
medication therapy	Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion				
if necessary	reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea,				
	cognitive changes, generalized rash, chest pain/pressure				
	Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate				
	hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors,				
	localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping,				
	uneasiness, agitation, feeling of impending doom				
	Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe White the second secon				
	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or				
	tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure				
Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hyperconsitivity (infusion reaction (grade 3) including systelia RD 90 00 mm I/g, bradycardia or					
	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure				
	tacifycardia, ffypoxerilia, dyspinea, cogiiit	live Citai	iges, generalized rash, chest pani, pressure		
	I				
Provider Name:		Provid	ler Signature:		
Attending Physician Name: Prov		Provid	ler NPI:		
Office Phone Number: Office		Office	Fax Number:		
(If ordering provider is an advanced practice practitioner)					
Note: This order is valid for 12 months from date of physician signature.					