

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116 Phone: 810-844-7373 Fax: 810-844-7366

Leuprolide (ELIGARD®)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: ___/___/___/

Referral Status:
New Referral

Dose or Frequency Change

Renewal

Patient Address: Patient Phone Numbo Date of Birth:/_ Weight:kg H		Memb Secon	ary Insurance: nber ID: ndary Insurance: nber ID:
Indication:	Diagnosis 0):		Labs Ibs required. Labs to be ordered by physician. Iher:
□ Leuprolide □ Leuprolide	e 7.5 mg subcutaneous every 4 weeks e 22.5 mg subcutaneous every 12 weeks e 30 mg subcutaneous every 16 weeks e 45 mg subcutaneous every 24 weeks		
Pre-Medications and Pre-Protocol (ordered at physician discretion)	 Acetaminophen 650 mg PO once Loratadine 10 mg PO once Diphenhydramine once 25 mg 50 mg PO IV Famotidine 20 mg IV once Hydrocortisone 100 mg IV once Methylprednisolone 125 mg IVP once Other:		Hydration LR Sodium Chloride 0.9% OthermL atmL/hr Before During Following



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116 Phone: 810-844-7373 Fax: 810-844-7366

Flushing Protocol (pre and post medication)	□ Sodium Chloride 0.9% □ 5 mL □ 10 mL as needed for line care	HeparinUnits/mLmL as needed for line care	
Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy as necessary	 Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg) Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches, generalized pain, back pain, abdominal cramping Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to moderate anaphylaxis for 3 doses Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure 		
Provider Name:		Provider Signature:	
Attending Physician Name:		Provider NPI:	
Office Phone Number:		Office Fax Number:	
(If ordering provider is an advanced practice practitioner) Note: This order is valid for 12 months from date of physician signature.			