

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116

Phone: 810-844-7373 Fax: 810-844-7366

Mepolizumab (NUCALA®)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:// Referral Status: □ New Referral □ Dose or Frequency Change □ Renewal				
Patient Address: Patient Phone Number Date of Birth:/_ Weight:kg H		Member Secondar	Insurance: ID: ry Insurance: ID:	
Diagnosis Diagnosis Code (ICD-10): Indication: Target start date:		Labs No labs required. Labs to be ordered by physician. □ Other:		
Hold and notify provi	der: if patient presents with acute uncontrolled	l asthma e	xacerbation OR signs/sy	mptoms of herpes zoster
☐ 100 mg ☐ 300 mg	NUCALA®) 100 mg/ml prefilled syringe aneous injection (into the upper arm, thigh, or	abdomen)	once. Repeat every 4 w	reeks.
Pre-Medications and Pre-Protocol (ordered at physician discretion)	□ Acetaminophen 650 mg PO once □ Loratadine 10 mg PO once □ Diphenhydramine once □ 25 mg □ 50 mg □ PO □ IV □ Famotidine 20 mg IV once □ Hydrocortisone 100 mg IV once □ Methylprednisolone 125 mg IVP once □ Other:		☐ LR ☐ Sodium Chloride 0.9% ☐ Other mL a	



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116 Phone: 810-844-7373

Fax: 810-844-7366

Flushing Protocol (pre and post	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL as needed for line care	☐ HeparinUnits/mLmL as needed for line care		
medication) Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy if necessary	 Acetaminophen 650 mg once as needed generalized pain, back pain, abdominal of Albuterol 2.5 mg /3 mL (0.083%) nebulized bronchospasm, wheezing, dyspnea, for 2 and Albuterol HFA inhaler 2 puffs q4hs as needed. Epinephrine injection 0.3 mg IM every 1 moderate anaphylaxis for 3 doses. Famotidine injection 20 mg IV over 2 min reaction including systolic BP 80-90 mmH cognitive changes, generalized rash, cheeded. Diphenhydramine injection 25 mg IV over hypersensitivity/infusion reaction (grade localized rash/hives, vomiting, nausea, puneasiness, agitation, feeling of impendial piphenhydramine injection 50 mg IV over hypersensitivity/infusion reaction (grade tachycardia, hypoxemia, dyspnea, cognition Hydrocortisone sodium succinate injection hypersensitivity/infusion reaction (grade) 	ter solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, 2 doses seeded for hypoxemia, bronchospasm, wheezing, dyspnea 5 minutes as needed for SBP less than 90mmHg, mild to nutes once as needed for severe hypersensitivity/infusion Hg, bradycardia or tachycardia, hypoxemia, dyspnea, st pain/pressure er 1 minute once as needed, moderate 2) including temperature greater than 100.5, rigors, bruritis, flushing, dizziness, back pain, abdominal cramping, and doom er 1 minute once as needed for severe 2 3), including systolic BP 80-90 mmHg, bradycardia or tive changes, generalized rash, chest pain/pressure		
Provider Name:		Provider Signature:		
Attending Physician Name:		Provider NPI:		
Office Phone Number:		Office Fax Number:		
(If ordering provider is an advanced practice practitioner) Note: This order is valid for 12 months from date of physician signature.				