

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116 Phone: 810-844-7373 Fax: 810-844-7366

Rabies Vaccine Follow up (Imovax[®], Rabavert[®])

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: ____/___/____

Referral Status:
New Referral
Dose or Frequency Change
Renewal

Patient Address: Patient Phone Numbe Date of Birth:/_ Weight:kg H		Memb Secon	ary Insurance: nber ID: ndary Insurance: nber ID:
Indication:	Diagnosis 0):	□ Otł	Labs
Rabies Vaccine Dose ⊠ 1ml IM □ Other:	Or X	ate of fir Day 4 Day 8 Day 15	dose considered day 1 irst dose:
Pre-Medications and Pre-Protocol (ordered at physician discretion) Flushing Protocol	 Acetaminophen 650 mg PO once Loratadine 10 mg PO once Diphenhydramine once 25 mg 50 mg PO IV Famotidine 20 mg IV once Hydrocortisone 100 mg IV once Methylprednisolone 125 mg IVP once Other: 		Hydration LR Sodium Chloride 0.9% OthermL atmL/hr Before During Following eparinUnits/mLmL
Flushing Protocol (pre and post medication)	□ Sodium Chloride 0.9% □ 5 mL □ 10 mL as needed for line care		eparinUnits/mLmL eeded for line care



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116 Phone: 810-844-7373 Fax: 810-844-7366

Hypersensitivity	• Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg)		
Panel	• Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches,		
Together Care	generalized pain, back pain, abdominal cramping		
Hypersensitivity Panel will be	• Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses		
ordered to provide emergency	 Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to 		
supportive care	moderate anaphylaxis for 3 doses		
medication therapy if necessary	 Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate 		
	 hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure 		
Provider Name:	Provider Signature:		
Attending Physician N	Name: Provider NPI:		
Office Phone Numbe	r: Office Fax Number:		
	s an advanced practice practitioner) lid for 12 months from date of physician signature.		