

### Immune Globulin IV (IGIV)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVE Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: \_\_\_/\_\_\_/\_\_\_

Referral Status:  New Referral  Dose or Frequency Change  Renewal

<b>Patient Name:</b> _____ <b>Patient Address:</b> _____ <b>Patient Phone Number:</b> _____ <b>Date of Birth:</b> ___/___/___ <b>Weight:</b> ___ kg <b>Height:</b> ___ cm <b>Allergies:</b> _____	<b>Primary Insurance:</b> _____ <b>Member ID:</b> _____ <b>Secondary Insurance:</b> _____ <b>Member ID:</b> _____	
<b>Diagnosis</b>	<b>Labs</b>	
Diagnosis Code (ICD-10): Other _____ Indication: _____ Target start date: _____	No labs required. Labs to be ordered by physician. <input type="checkbox"/> Other: _____	
<b>Immune Globulin Intravenous (Gamunex C Liquid 10%, Octagam 10% Liquid or Gammagard Liquid 10% or other IVIG products based on availability) via IV infusion per ArrIVE Standards for the Prescribing and Administration of Immune Globulin in Adult Patients. ___ gram/kg* _____ grams</b>		
<b>Frequency:</b> <input type="checkbox"/> Once <input type="checkbox"/> Daily <input type="checkbox"/> Every _____ week(s) <input type="checkbox"/> Other _____	<b>Duration:</b> <input type="checkbox"/> _____ Doses <input type="checkbox"/> Other _____	
<b>Special orders:</b> _____ (i.e., repeat every 12 weeks)		
* Dose will be calculated based on IBW or adjusted body weight as applicable, and rounded per Together Care dose-rounding logic		
<b>Pre-Medications and Pre-Protocol</b> (ordered at physician discretion)	<input type="checkbox"/> Acetaminophen 650 mg PO once <input type="checkbox"/> Loratadine 10 mg PO once <input type="checkbox"/> Diphenhydramine once <input type="checkbox"/> 25 mg <input type="checkbox"/> 50 mg <input type="checkbox"/> PO <input type="checkbox"/> IV <input type="checkbox"/> Famotidine 20 mg IV once <input type="checkbox"/> Hydrocortisone 100 mg IV once <input type="checkbox"/> Methylprednisolone 125 mg IVP once <input type="checkbox"/> Other: _____	<b>Hydration</b> <input type="checkbox"/> LR <input type="checkbox"/> Sodium Chloride 0.9% <input type="checkbox"/> Other _____  _____ mL at _____ mL/hr <input type="checkbox"/> Before <input type="checkbox"/> During <input type="checkbox"/> Following

<b>Flushing Protocol</b> <i>as needed for line care</i>	<input type="checkbox"/> <b>Sodium Chloride 0.9%</b> <input type="checkbox"/> 5 mL <input type="checkbox"/> 10 mL	<input type="checkbox"/> <b>Heparin 100 Units/mL</b> <input type="checkbox"/> 5 mL <input type="checkbox"/> 10 mL
<b>Hypersensitivity Panel</b> Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy if necessary	<ul style="list-style-type: none"> <li>• <b>Sodium chloride</b> 0.9% bolus 500 mL once as needed for hypotensive management (SBP &lt;90mmHg)</li> <li>• <b>Acetaminophen</b> 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches, generalized pain, back pain, abdominal cramping</li> <li>• <b>Albuterol</b> 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses</li> <li>• <b>Albuterol HFA</b> inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea</li> <li>• <b>Epinephrine</b> injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to moderate anaphylaxis for 3 doses</li> <li>• <b>Famotidine</b> injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure</li> <li>• <b>Diphenhydramine</b> injection 25 mg IV over 1 minute once as needed, moderate hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom</li> <li>• <b>Diphenhydramine</b> injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure</li> <li>• <b>Hydrocortisone</b> sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure</li> </ul>	
Provider Name: _____ Provider Signature: _____ Attending Physician Name: _____ Provider NPI: _____ Office Phone Number: _____ Office Fax Number: _____ <i>Note: This order is valid for 12 months from date of physician signature.</i>		