

An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116-2084

Phone: 810-844-7373 Fax: 810-844-7366

## Denosumab (XGEVA or Biosimilar)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:// Referral Status: □ New Referral □ Dose or Frequency Change □ Renewal				
Patient Address: Patient Phone Numb Date of Birth:/_ Weight:kg		Member ID: Secondary Insurance:		
Indication:	Diagnosis 10):	Labs  Albumin  Magnesium Creatinine (serum) Calcium Other:	$\square$ Prior to each injection	
NOTE TO PROVIDER: All patients with Denosumab prescribed should receive at least 1000 mg Calcium and 400 IU Vitamin D daily per prescribing information.  Hold and notify provider: Notify provider and hold at provider discretion for Ca < 7 mg/dL or Magnesium < 1.5 mg/dL. Calcium and magnesium level should be corrected prior to initiation of treatment.				
Denosumab (Xgeva or Biosimilar) 120 mg subcutaneous injection  Frequency:				
Pre-Medications and Pre-Protocol (ordered at physician discretion)	□ Acetaminophen 650 mg PO once     □ Loratadine 10 mg PO once     □ Diphenhydramine once     □ 25 mg □ 50 mg     □ PO □ IV     □ Famotidine 20 mg IV once     □ Hydrocortisone 100 mg IV once     □ Methylprednisolone 125 mg IVP once     □ Other:	☐ LR ☐ Sodium Chloride 0. ☐ Other		



An Outpatient Department of Trinity Health Oakland 9400 Village Place Blvd, Brighton, MI 48116-2084

Phone: 810-844-7373 Fax: 810-844-7366

Flushing Protocol	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL	☐ HeparinUnits/mLmL		
(pre and post	as needed for line care	as needed for line care		
medication)				
Hypersensitivity	• Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg)			
Panel	• Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches,			
Together Care	generalized pain, back pain, abdominal cramping			
Hypersensitivity	Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia,			
Panel will be	bronchospasm, wheezing, dyspnea, for 2 doses			
ordered to provide	<ul> <li>Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea</li> </ul>			
emergency	• Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to			
supportive care	moderate anaphylaxis for 3 doses			
medication	• Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion			
therapy as	reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive			
necessary	changes, generalized rash, chest pain/pressure			
	Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate			
	hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized			
		rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness,		
		agitation, feeling of impending doom		
		Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe		
	1 1 1	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or		
		tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure		
	The state of the s	Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe		
		hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or		
	tachycardia, hypoxemia, dyspnea, cognitive cha	tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure		
Provider Name:		Provider Signature:		
Trovider Name:		Trovider signature.		
Attending Physician Name:		Provider NPI:		
<b>0</b> ,				
Office Phone Numbe	er:	Office Fax Number:		
(If ordering provider is an advanced practice practitioner)				
Note: This order is valid for 12 months from date of physician signature.				