ArrIVe
INFUSION THERAPY

An Outpatient Department of Trinity Health Oakland
9400 Village Place Blvd, Brighton, MI 48116-2084
Phone: 810-844-7373
Fax: 810-844-7366

Ocrelizumab and Hyaluronidase (OCREVUS ZUNOVO) or Biosimilar Order Form

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: ____/____/____

Referral Status: ☐ New Referral ☐ Dose or Frequency Change ☐ Renewal

Patient Name: _____ **Patient Address:** _____ **Patient Phone Number:** _____ **Date of Birth**: ____/____/____ **Weight:** _____kg **Height:** _____cm **Allergies:** _____	**Primary Insurance:** _____ **Member ID:** _____ **Secondary Insurance:** _____ **Member ID:** _____
Diagnosis Diagnosis Code (ICD-10): Other _____ Indication: _____ Target start date: _____	**Labs** **Lab orders (within 45 days):** Prior to start of therapy: Hepatitis B surface antigen, Hepatitis B surface antibody, Hepatitis B core antibody IgM Every 26 weeks (6 months): Complete blood count and differential, hepatic function panel, Immunoglobulin IgM, and Immunoglobulin IgG ☐ Other: _____ **Hold and notify physician:** - Abnormal hepatitis B and TB test results - Signs or symptoms of active infection/fever prior to infusion/treatment

Ocrevus and Hyaluronidase (OCREVUS ZUNOVO) or Biosimilar 920mg subcutaneous injection over 10 minutes.

Nursing Orders:
Monitor for signs/symptoms of injection reaction during administration and for at least 1 hour following first injection; monitor for at least 15 minutes following subsequent injections.

Frequency:
☐ Once ☐ Daily ☐ Every _____ week(s)
☐ Other ☒ 6 months_____

ArrIVe
INFUSION THERAPY

An Outpatient Department of Trinity Health Oakland
9400 Village Place Blvd, Brighton, MI 48116-2084
Phone: 810-844-7373
Fax: 810-844-7366

Pre-Medications and Pre-Protocol (ordered at physician discretion)	☐ Acetaminophen 650 mg PO once ☐ Loratadine 10 mg PO once ☒ Diphenhydramine 30-60 min prior to infusion ☒25 mg ☐50 mg ☒ PO ☐IV ☐ Famotidine 20 mg IV once ☐ Hydrocortisone 100 mg IV once ☒ Dexamethasone 20 mg PO 30-60 min prior to infusion ☐ Other: _____	**Hydration** ☐ LR ☐ Sodium Chloride 0.9% ☐ Other _____ _____ mL at _____mL/hr ☐ Before ☐ During ☐ Following

ArrIVe INFUSION THERAPY

An Outpatient Department of Trinity Health Oakland
9400 Village Place Blvd, Brighton, MI 48116-2084
Phone: 810-844-7373
Fax: 810-844-7366

Flushing Protocol (pre and post medication)	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL as needed for line care	☐ Heparin 100 Units/mL ☐ 5 mL ☐ 10 mL as needed for line care
Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy if necessary	• **Sodium chloride** 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg) • **Acetaminophen** 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches, generalized pain, back pain, abdominal cramping • **Albuterol** 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses • **Albuterol HFA** inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea • **Epinephrine** injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to moderate anaphylaxis for 3 doses • **Famotidine** injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure • **Diphenhydramine** injection 25 mg IV over 1 minute once as needed, moderate hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom • **Diphenhydramine** injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure • **Hydrocortisone** sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure	

Provider Name: _____ Provider Signature: _____

Attending Physician Name: _____ Provider NPI: _____

Office Phone Number: _____ Office Fax Number: _____

(If ordering provider is an advanced practice practitioner)
Note: This order is valid for 12 months from date of physician signature.