ArrIVe INFUSION THERAPY

An Outpatient Department of St. Mary's Hospital
2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325
Phone: 706-389-7802
Fax: 706-389-2501

Belimumab (Benlysta®)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: ____/____/____

Referral Status: ☐ New Referral ☐ Dose or Frequency Change ☐ Renewal

Patient Name: _____ **Patient Address:** _____ **Patient Phone Number:** _____ **Date of Birth**: ____/____/____ **Weight:** _____kg **Height:** _____cm **Allergies:** _____	**Primary Insurance:** _____ **Member ID:** _____ **Secondary Insurance:** _____ **Member ID:** _____

Diagnosis	**Labs**	**Frequency**
☐ Drug-induced systemic lupus erythematosus (M32.0) ☐ Systemic lupus erythematosus organ or system involvement unspecified (M32.10) ☐ Endocarditis in systemic lupus erythematosus (M32.11) ☐ Pericarditis in systemic lupus erythematosus (M32.12) ☐ Lung involvement in systemic lupus erythematosus (M32.13) ☐ Glomerular disease in systemic lupus erythematosus (M32.14) ☐ Tubulo-interstitial nephropathy in systemic lupus erythematosus (M32.15) ☐ Other organ or system involvement in systemic lupus erythematosus (M32.19) ☐ Other forms of systemic lupus erythematosus (M32.8) ☐ Systemic lupus erythematosus, unspecified (M32.9) ☐ Other Diagnosis Code (ICD-10): _____ ☐ Other Indication: _____ Target start date: _____	☐ CMP ☐ CBC w/diff ☐ CBC w/o diff	☐ Every infusion ☐ Other: _____

Previously tried and failed therapies (include dates): _____

Belimumab (Benlysta)

Dose

☐ 10 mg/kg
☐ _____ mg/kg
☐ _____ mg

Frequency

Induction	**Maintenance**
☐ On weeks 0, 2, 4	☐ Every 4 weeks ☐ Every _____ weeks

Date of last infusion: ____/____/____

ArrIVe INFUSION THERAPY

An Outpatient Department of St. Mary's Hospital
2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325
Phone: 706-389-7802
Fax: 706-389-2501

Pre-Medications and Pre-Protocol (ordered at physician discretion)	☐ Acetaminophen 650 mg PO once ☐ Loratadine 10 mg PO once ☐ Diphenhydramine once ☐25 mg ☐50 mg ☐ PO ☐IV ☐ Famotidine 20 mg IV once ☐ Hydrocortisone 100 mg IV once ☐ Methylprednisolone 125 mg IVP once ☐ Other: _____	**Hydration** ☐ LR ☐ Sodium Chloride 0.9% ☐ Other _____ _____ mL at _____mL/hr ☐ Before ☐ During ☐ Following
Flushing Protocol *as needed for line care*	☐Sodium Chloride 0.9% ☐5 mL☐10 mL	☐Heparin 100 Units/mL☐5 mL☐10 mL
Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy as necessary.	• **Sodium chloride** 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg) • **Acetaminophen** 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches, generalized pain, back pain, abdominal cramping • **Albuterol** 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses • **Albuterol HFA** inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea • **Epinephrine** injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to moderate anaphylaxis for 3 doses • **Famotidine** injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure • **Diphenhydramine** injection 25 mg IV over 1 minute once as needed, moderate hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom • **Diphenhydramine** injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure • **Hydrocortisone** sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure	

Provider Name: _____ Provider Signature: _____

Attending Physician Name: _____ Provider NPI: _____

Office Phone Number: _____ Office Fax Number: _____

Note: This order is valid for 12 months from date of physician signature.

Submit