

Denosumab Biosimilar

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVE Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: ___/___/___

Referral Status: New Referral Dose or Frequency Change Renewal

Patient Name: _____ Patient Address: _____ Patient Phone Number: _____ Date of Birth: ___/___/___ Weight: ___ kg Height: ___ cm Allergies: _____	Primary Insurance: _____ Member ID: _____ Secondary Insurance: _____ Member ID: _____	
<p style="text-align: center;">Diagnosis</p> Diagnosis Code (ICD-10): _____ Indication: _____ Target start date: _____	<p style="text-align: center;">Labs</p> <input type="checkbox"/> Albumin <input type="checkbox"/> Magnesium <input type="checkbox"/> Once <input type="checkbox"/> Creatinine (serum) <input type="checkbox"/> Monthly <input type="checkbox"/> Calcium <input type="checkbox"/> Prior to each injection <input type="checkbox"/> Other: _____ <input type="checkbox"/> Other: _____	
<p>NOTE TO PROVIDER: All patients with Denosumab (Xgeva®) prescribed should receive at least 1000 mg Calcium and 400 IU Vitamin D daily per prescribing information.</p>		
<p>Hold and notify provider: Notify provider and hold at provider discretion for Ca < 7 mg/dL or Magnesium < 1.5 mg/dL. Calcium and magnesium level should be corrected prior to initiation of treatment.</p>		
<p>Denosumab 120 mg subcutaneous injection</p> <p style="text-align: center;">Frequency: _____</p> <p>(Bomynta, Osenvelt, Wyost, Xgeva or other biosimilars)</p>		
Pre-Medications and Pre-Protocol (ordered at physician discretion)	<input type="checkbox"/> Acetaminophen 650 mg PO once <input type="checkbox"/> Loratadine 10 mg PO once <input type="checkbox"/> Diphenhydramine once <input type="checkbox"/> 25 mg <input type="checkbox"/> 50 mg <input type="checkbox"/> PO <input type="checkbox"/> IV <input type="checkbox"/> Famotidine 20 mg IV once <input type="checkbox"/> Hydrocortisone 100 mg IV once <input type="checkbox"/> Methylprednisolone 125 mg IVP once <input type="checkbox"/> Other: _____	<p style="text-align: center;">Hydration</p> <input type="checkbox"/> LR <input type="checkbox"/> Sodium Chloride 0.9% <input type="checkbox"/> Other: _____ _____ mL at _____ mL/hr <input type="checkbox"/> Before <input type="checkbox"/> During <input type="checkbox"/> Following

Flushing Protocol <i>as needed for line care</i>	<input type="checkbox"/> Sodium Chloride 0.9% <input type="checkbox"/> 5 mL <input type="checkbox"/> 10 mL	<input type="checkbox"/> Heparin 100 Units/mL <input type="checkbox"/> 5 mL <input type="checkbox"/> 10 mL
Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy as necessary	<ul style="list-style-type: none"> Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg) Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches, generalized pain, back pain, abdominal cramping Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to moderate anaphylaxis for 3 doses Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure 	
Provider Name: _____		Provider Signature: _____
Attending Physician Name: _____		Provider NPI: _____
Office Phone Number: _____		Office Fax Number: _____
<i>Note: This order is valid for 12 months from date of physician signature.</i>		