

An Outpatient Department of St. Mary's Hospital
2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325
Phone: 706-389-7802
Fax: 706-389-2501

Rabies Vaccine Follow up (Imovax®, Rabavert®)

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVE Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date: ____/____/____

Referral Status: ☐ New Referral ☐ Dose or Frequency Change ☐ Renewal

Patient Name: _____ Patient Address: _____ Patient Phone Number: _____ Date of Birth: ____/____/____ Weight: ____ kg Height: ____ cm Allergies: _____		Primary Insurance: _____ Member ID: _____ Secondary Insurance: _____ Member ID: _____	
Diagnosis Diagnosis Code (ICD-10): _____ Indication: _____ Target start date: _____		Labs ☐ Other: _____	
Rabies Vaccine Intramuscular Injection Dose ☒ 1ml IM ☐ Other: _____ Frequency Original dose considered day 1 Date of first dose: _____ ☒ Day 4 ☒ Day 8 ☒ Day 15 ☐ Other: _____			
Pre-Medications and Pre-Protocol (ordered at physician discretion)		Hydration ☐ LR ☐ Sodium Chloride 0.9% ☐ Other: _____ _____ mL at _____ mL/hr ☐ Before ☐ During ☐ Following	
Flushing Protocol <i>as needed for line care</i>		☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL ☐ Heparin 100 Units/mL ☐ 5 mL ☐ 10 mL	

Hypersensitivity Panel Together Care Hypersensitivity Panel will be ordered to provide emergency supportive care medication therapy if necessary	• Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg) • Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches, generalized pain, back pain, abdominal cramping • Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses • Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea • Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to moderate anaphylaxis for 3 doses • Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure • Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom • Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure • Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure
Provider Name: _____ Provider Signature: _____ Attending Physician Name: _____ Provider NPI: _____ Office Phone Number: _____ Office Fax Number: _____ <i>Note: This order is valid for 12 months from date of physician signature.</i>	