

An Outpatient Department of St. Mary's Hospital 2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325

Phone: 706-389-7802 Fax: 706-389-2501

Rituximab (Rituxan®) or Biosimilar

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:// Referral Status: □ New Referral □ Dose or Frequency Change □ Renewal	
Patient Name: Patient Address: Patient Phone Number: Date of Birth: Weight: kg Height: Allergies:	Primary Insurance: Member ID: Secondary Insurance: Member ID:
Diagnosis Diagnosis Code (ICD-10): Indication: Target start date: Note to provider: Viral hepatitis B screening required prior to the	Labs CBC w/ diff (specify frequency): Other: rapy initiation. Additional screening for hepatitis C, HIV, and TB
may be warranted. Hold and Notify Provider: ANC below 1.5, Plt below 75K; signs/sy	mptoms of active infection.
Rituximab (Or Biosimilar) Pharmacy to Select Truxima (rituximab-abbs) Ruxience (rituximab-pvvr) Rituxan (rituximab)	
Dose ☐ 1000 mg IVPB ☐ 375 mg/m² IVPB ☐ 500 mg IVPB ☐ Other:	Frequency ☐ Day 1 and 15, ☐ Repeating every 6 months ☐ Weekly for weeks ☐ Once ☐ Other: Note: interval to be no less than 20 weeks from day 1 dose of previous cycle



An Outpatient Department of St. Mary's Hospital 2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325 Phone: 706-389-7802 Fax: 706-389-2501

Pre-Medications and Pre-Protocol (ordered at physician discretion)	△ Acetaminophen 650 mg PO once ☐ Loratadine 10 mg PO once ☑ Diphenhydramine once 25 mg	Hydration LR Sodium Chloride 0.9% OthermL atmL/hr Before During Following
Flushing Protocol as needed for line care	Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL	Heparin 100 Units/mL □ 5 mL □ 10 mL
 Hypersensitivity Panel Together Care Hypersensitivity Panel volume of the provide emergency Supportive care medication therapy as necessary Albuterol 2.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea, for 2 doses Albuterol 4.5 mg /3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia, bronchospasm, wheezing, dyspnea emedication therapy as necessary Finephrine injection 0.3 mg IM every 15 minutes as needed for severe hypersensitivity/infusion reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Diphenhydramine injection 25 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors, localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping, uneasiness, agitation, feeling of impending doom Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure 		
Provider Name:		Provider Signature:
Attending Physician N	ame:	Provider NPI:
Office Phone Number		Office Fax Number:
(If ordering provider is an advanced practice practitioner) Note: This order is valid for 12 months from date of physician signature.		