

An Outpatient Department of St. Mary's Hospital 2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325

Phone: 706-389-7802 Fax: 706-389-2501

Tocilizumab (Actemra ®) or Biosimilar

Along with this order form, please include the following: Demographics, Insurance Information, Lab Results, Current Medications, and Recent Visit Notes. ArrIVe Infusion will obtain any necessary medication authorizations for patients receiving infusion therapies.

Order Date:// Referral Status: □ New Referral □ Dose or Frequency Change □ Renewal			
Patient Name:	Primary Insurance:		
Patient Address:	Member ID:	Member ID:	
Patient Phone Number:	Secondary Insurance:		
Date of Birth:/	Member ID:	Member ID:	
Weight:kg Height:cm			
Allergies:			
Diagnosis	Labs	Frequency	
☐ Rheumatoid arthritis with rheumatoid factor, unspecified	☐ LFTs		
(M05.9)	☐ CBC w/ diff	\square Every infusion	
☐ Rheumatoid arthritis without rheumatoid factor, other specifie	ed ☐ CBC w/o diff	☐ Other:	
site (M06.0A)	☐ Lipid panel		
☐ Rheumatoid arthritis, unspecified (M06.9)	☐ Other:		
\square Juvenile rheumatoid arthritis with systemic onset, other specified site (M08.2A)			
☐ Juvenile rheumatoid polyarthritis (seronegative) (M08.3)			
☐ Giant cell arthritis with polymyalgia rheumatica (M31.5)			
☐ Other giant cell arthritis (M31.6)			
☐ Systemic sclerosis with lung involvement (M34.81)			
☐ Other Diagnosis Code (ICD-10):			
☐ Other Indication:			
Target start date:			
Previously tried and failed therapies (include dates):			
Tocilizumab (Actemra ®) or Biosimilar in 100 mL of 0.9% sodium chloride or 50 mL of 0.9 sodium chloride for <30 kg Nursing note: Allow the fully diluted solution to reach room temperature prior to infusion. Protect from light.			
Dose	Frequency		
☐ 4 mg/kg	☐ Every 4 weeks		
☐ 6 mg/kg	□ Every 4 weeks		
□ 8 mg/kg	Livery weeks		
□mg/kg □mg	Date of last infusion:/	<i>J</i>	



An Outpatient Department of St. Mary's Hospital 2421 Malcom Bridge Rd Ste 820 Bogart, GA 30622-2325 Phone: 706-389-7802 Fax: 706-389-2501

Pre-Medications	☐ Acetaminophen 650 mg PO once	Hydration		
and Pre-Protocol	☐ Loratadine 10 mg PO once	□LR		
(ordered at	☐ Diphenhydramine once	☐ Sodium Chloride 0.9%		
physician discretion)	□25 mg □50 mg	☐ Other		
	□ PO □IV			
	☐ Famotidine 20 mg IV once	mL atmL/hr		
	☐ Hydrocortisone 100 mg IV once	☐ Before ☐ During ☐ Following		
	☐ Methylprednisolone 125 mg IVP once			
	☐ Other:			
Flushing Protocol	☐ Sodium Chloride 0.9% ☐ 5 mL ☐ 10 mL	Heparin 100 Units/L ☐ 5 mL ☐ 10 mL		
as needed for line care				
Hypersensitivity	• Sodium chloride 0.9% bolus 500 mL once as needed for hypotensive management (SBP <90mmHg)			
Panel	• Acetaminophen 650 mg once as needed for temperature GREATER than 38 C (100.4 F), headaches,			
Together Care	generalized pain, back pain, abdominal cramping			
Hypersensitivity	• Albuterol 2.5 mg/3 mL (0.083%) nebulizer solution 2.5 mg, nebulize every 10 min PRN, hypoxemia,			
Panel will be	bronchospasm, wheezing, dyspnea, for 2 doses			
ordered to provide	Albuterol HFA inhaler 2 puffs q4hs as needed for hypoxemia, bronchospasm, wheezing, dyspnea			
emergency	Epinephrine injection 0.3 mg IM every 15 minutes as needed for SBP less than 90mmHg, mild to			
supportive care	moderate anaphylaxis for 3 doses			
medication therapy	• Famotidine injection 20 mg IV over 2 minutes once as needed for severe hypersensitivity/infusion			
as necessary	reaction including systolic BP 80-90 mmHg, bradycardia or tachycardia, hypoxemia, dyspnea,			
	cognitive changes, generalized rash, chest pain/pressure			
	Diphenhydramine injection 25 mg IV over 1 minute once as needed, moderate			
	hypersensitivity/infusion reaction (grade 2) including temperature greater than 100.5, rigors,			
	localized rash/hives, vomiting, nausea, pruritis, flushing, dizziness, back pain, abdominal cramping,			
	 uneasiness, agitation, feeling of impending doom Diphenhydramine injection 50 mg IV over 1 minute once as needed for severe 			
	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or			
	tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure			
	 Hydrocortisone sodium succinate injection 100 mg IV once as needed for severe 			
	hypersensitivity/infusion reaction (grade 3), including systolic BP 80-90 mmHg, bradycardia or			
tachycardia, hypoxemia, dyspnea, cognitive changes, generalized rash, chest pain/pressure				
	, , , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,		
Provider Name:		Provider Signature:		
Attending Physician N	ame:	Provider NPI:		
Office Phone Number	:	Office Fax Number:		
Note: This order is valid for 12 months from date of physician				
	ia for 12 months from date of physician			
signature.				